



A service of the National Library of Medicine  
and the National Institutes of Health

My NCBI  
[Sign In] [Regis]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Bool

Search PubMed for [ ] Go Clear

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort by Send to

All: 1 Review: 0

1: Am J Pathol. 2006 Sep;169(3):806-14.

Full Text  
Am J Pathol

FREE full text article  
in PubMed Central

Links

**Odontogenic keratocysts arise from quiescent epithelial rests and are associated with deregulated hedgehog signaling in mice and humans.**

**Grachtchouk M, Liu J, Wang A, Wei L, Bichakjian CK, Garlick J, Paulino AF, Giordano T, Dlugosz AA.**

Department of Dermatology and Comprehensive Cancer Center, University of Michigan, 3316 CCGC, Box 0932, 1500 E. Medical Center Dr., Ann Arbor, MI 48109-0932, USA.

Odontogenic keratocysts in humans are aggressive, noninflammatory jaw cysts that may harbor PTCH1 mutations, leading to constitutive activity of the embryonic Hedgehog (Hh) signaling pathway. We show here that epithelial expression of the Hh transcriptional effector Gli2 is sufficient for highly penetrant keratocyst development in transgenic mice. Mouse and human keratocysts expressed similar markers, leading to tooth misalignment, bone remodeling, and craniofacial abnormalities. We detected Hh target gene expression in epithelial cells lining keratocysts from both species, implicating deregulated Hh signaling in their development. Most mouse keratocysts arose from rests of Malassez--quiescent, residual embryonic epithelial cells that remain embedded in the periodontal ligament surrounding mature teeth. In Gli2-expressing mice, these rests were stimulated to proliferate, stratify, and form a differentiated squamous epithelium. The frequent development of keratocysts in Gli2-expressing mice supports the idea that GLI transcription factor activity mediates pathological responses to deregulated Hh signaling in humans. Moreover, Gli2-mediated reactivation of quiescent epithelial rests to form keratocysts indicates that these cells retain the capacity to function as progenitor cells on activation by an appropriate developmental signal.

PMID: 16936257 [PubMed - indexed for MEDLINE]

#### Related Links

Sustained Hedgehog signaling is required for basal cell carcinoma proliferation and survival: conditional skin tumorigenesis recapitulates the hair cycle. [Cancer Res. 2005]

A previously unidentified amino-terminal domain regulates transcriptional activity of wild-type and disease-associated Gli3. [Cancer Res. 2005]

Functional compensation in Hedgehog signaling during mouse prostate development. [Dev Biol. 2006]

Hedgehog pathway activity in the LADY prostate tumor model. [Cancer Res. 2007]

Unique and complimentary activities of the Gli transcription factors in Hedgehog signaling. [Exp Cell Res. 2006]

See all Related Articles...

Exhibit 2

Display AbstractPlus Show 20 Sort by Send to

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services